Cambridge IGCSE

Cambridge Assessment International Education
Cambridge International General Certificate of Secondary Education

CANDIDATE NAME

CENTRE NUMBER | | | | | |

CANDIDATE NUMBER | | | | |

3629900230

ADDITIONAL MATHEMATICS **0606/23**

Paper 2 **October/November 2019**

2 hours

Candidates answer on the Question Paper.

Additional Materials: Electronic calculator

READ THESE INSTRUCTIONS FIRST

Write your centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 80.

This document consists of **15** printed pages and **1** blank page.

© UCLES 2019

Cambridge Assessment International Education

[Turn over

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

Binomial Theorem

$$(a + b)^n = a^n + \binom{n}{1} a^{n-1} b + \binom{n}{2} a^{n-2} b^2 + \ldots + \binom{n}{r} a^{n-r} b^r + \ldots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$.

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2} bc \sin A$$

© UCLES 2019

1 Solve $|3x+2| = x+4$. [3]

2 **(i)** Show that $\dfrac{\text{cosec}x - \cot x}{1-\cos x} = \text{cosec}x$. [3]

(ii) Hence solve $\dfrac{\text{cosec}x - \cot x}{1-\cos x} = 2$ for $0° < x < 180°$. [2]

3 The first four terms in the expansion of $(1+ax)^5(2+bx)$ are $2+32x+210x^2+cx^3$, where a, b and c are integers. Show that $3a^2-16a+21=0$ and hence find the values of a, b and c. [8]

© UCLES 2019

4 **(i)** Given that $y = 2x^2 - 4x - 7$, write y in the form $a(x-b)^2 + c$, where a, b and c are constants. [3]

(ii) Hence write down the minimum value of y and the value of x at which it occurs. [2]

(iii) Using your answer to **part (i)**, solve the equation $2p - 4\sqrt{p} - 7 = 0$, giving your answer correct to 2 decimal places. [3]

© UCLES 2019

5 **(a)** Solve $3\cot^2\left(y-\frac{\pi}{4}\right)=1$ for $0<y<\pi$ radians. [4]

(b) Solve $7\cot z+\tan z=7\operatorname{cosec} z$ for $0°\leqslant z\leqslant 360°$. [6]

© UCLES 2019

6 Do not use a calculator in this question.



(i) Find tan ACB in the form $r + s\sqrt{3}$, where r and s are integers. [3]

(ii) Find AC in the form $t\sqrt{u}$, where t and u are integers and $t \neq 1$. [3]

7



The diagram shows part of the curve $y = x + \frac{6}{(3x+2)^2}$ and the line $x = 2$.

(i) Find, correct to 2 decimal places, the coordinates of the stationary point. [6]

© UCLES 2019

(ii) Find the area of the shaded region, showing all your working. [4]

8 The roots of the equation $x^3 + ax^2 + bx + 24 = 0$ are 2, 3 and p, where p is an integer.

(i) Find the value of p. [1]

(ii) Show that $a = -1$ and find the value of b. [4]

© UCLES 2019

Given that a curve has equation $y = x^3 - x^2 + bx + 24$ find, using your value of b,

(iii) $\frac{\mathrm{d}y}{\mathrm{d}x}$, [1]

(iv) the integer value of x for which the gradient of the curve is 2 and the corresponding value of y. [3]

The coordinates of the point P on the curve are given by the values of x and y found in **part (iv)**.

(v) Find the equation of the tangent to the curve at P. [1]

© UCLES 2019

9



The diagram shows points O, A, B, C, D and X. The position vectors of A, B, and C relative to O are $\overrightarrow{OA} = \mathbf{a}$, $\overrightarrow{OB} = 2\mathbf{b}$ and $\overrightarrow{OC} = 3\mathbf{a}$. The vector $\overrightarrow{CD} = \mathbf{b}$.

(i) Given that $\overrightarrow{AX} = \lambda \overrightarrow{AD}$, find $\overrightarrow{OX}$ in terms of λ, **a** and **b**. [2]

(ii) Given that $\overrightarrow{BX} = \mu \overrightarrow{BC}$, find $\overrightarrow{OX}$ in terms of μ, **a** and **b**. [2]

© UCLES 2019

(iii) Hence find the value of λ and of μ. [4]

(iv) Find the ratio $\frac{AX}{XD}$. [1]

10 The functions f and g are defined by

$$\mathrm{f}(x) = \ln(3x+2) \quad \text{for } x > -\frac{2}{3},$$
$$\mathrm{g}(x) = \mathrm{e}^{2x} - 4 \quad \text{for } x \in \mathbb{R}.$$

(i) Solve $\mathrm{gf}(x) = 5$. [5]

© UCLES 2019

(ii) Find $f^{-1}(x)$. [2]

(iii) Solve $f^{-1}(x) = g(x)$. [4]

© UCLES 2019